



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON CERTAIN THEOREMS OF JACOBI RELATING TO CANONICAL EQUATIONS.

By PROF. ORMOND STONE, Charlottesville, Va.

Jacobi in his Vorlesungen über Dynamik first proves the particular theorem contained in Art. 5 of the present paper, and afterwards derives therefrom that given in Art. 6. In what follows it will be seen that each of these is made a special case of the more general theorem given in Art. 1.

1. If we have given the partial differential equations

$$\frac{\partial S}{\partial t} + H = 0, \quad (\text{a})$$

$$y_1 = \frac{\partial S}{\partial x_1}, \quad y_2 = \frac{\partial S}{\partial x_2}, \quad \dots, \quad y_{2n} = \frac{\partial S}{\partial x_{2n}}, \quad (\text{b})$$

in which H is a function of t and of the x 's and y 's, but S is a function of t and the x 's only ; and if there exist the so-called canonical relations

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial y_i}, \quad \frac{dy_i}{dt} = -\frac{\partial H}{\partial x_i}; \quad (i = 1, 2, \dots, n) \quad (c)$$

then will similar canonical relations exist between the remaining variables ; that is to say, we shall have

$$\frac{dx_{n+i}}{dt} = \frac{\partial H}{\partial y_{n+i}}, \quad \frac{dy_{n+i}}{dt} = -\frac{\partial H}{\partial x_{n+i}}. \quad (i = 1, 2, \dots, n) \quad (d)$$

Differentiating each of the y 's totally with reference to t , we have

$$\left. \begin{aligned} \frac{\partial y_1}{\partial t} + \sum_1^{2n} \frac{\partial y_1}{\partial x_i} \frac{dx_i}{dt} - \frac{dy_1}{dt} &= 0, \\ \frac{\partial y_2}{\partial t} + \sum_1^{2n} \frac{\partial y_2}{\partial x_i} \frac{dx_i}{dt} - \frac{dy_2}{dt} &= 0, \\ . &. \\ \frac{\partial y_{2n}}{\partial t} + \sum_1^{2n} \frac{\partial y_{2n}}{\partial x_i} \frac{dx_i}{dt} - \frac{dy_{2n}}{dt} &= 0. \end{aligned} \right\} \quad (1)$$

Also, since

$$\frac{\partial}{\partial x_i} \frac{\partial S}{\partial t} = \frac{\partial}{\partial t} \frac{\partial S}{\partial x_i} = \frac{\partial y_i}{\partial t},$$

if we differentiate equations (a) partially with reference to x_1, x_2, \dots, x_{2n} , we shall have

$$\left. \begin{aligned} \frac{\partial y_1}{\partial t} + \frac{\partial H}{\partial x_1} + \sum_1^{2n} \frac{\partial H}{\partial y_i} \frac{\partial y_i}{\partial x_1} &= 0, \\ \frac{\partial y_2}{\partial t} + \frac{\partial H}{\partial x_2} + \sum_1^{2n} \frac{\partial H}{\partial y_i} \frac{\partial y_i}{\partial x_2} &= 0, \\ . &. \\ \frac{\partial y_{2n}}{\partial t} + \frac{\partial H}{\partial x_{2n}} + \sum_1^{2n} \frac{\partial H}{\partial y_i} \frac{\partial y_i}{\partial x_{2n}} &= 0. \end{aligned} \right\} \quad (2)$$

Since

$$\frac{\partial}{\partial x_i} \frac{\partial S}{\partial x_i} = \frac{\partial}{\partial x_i} \frac{\partial S}{\partial x_i},$$

we may write

$$\frac{\partial y_i}{\partial x_j} = \frac{\partial y_j}{\partial x_i} = a_{i,j};$$

whence by reason of equations (c), we may also write

[illegible]

same ; hence the corresponding unknown quantities must be the same : whence

$$\frac{dx_{n+i}}{dt} = \frac{\partial H}{\partial y_{n+i}}, \quad \frac{dy_{n+i}}{dt} = -\frac{\partial H}{\partial x_{n+i}}. \quad (i = 1, 2, \dots, n) \quad (d)$$

Q. E. D.

2. It may be proved similarly that if

$$\frac{dx_{n+1}}{dt} + \frac{\partial H}{\partial y_{n+1}}, \quad \frac{dy_{n+1}}{dt} = -\frac{\partial H}{\partial x_{n+1}};$$

then

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial y_i}, \quad \frac{dy_i}{dt} = -\frac{\partial H}{\partial x_i},$$

and, in general, if any $2n$ of the $4n$ relations (c) and (d) exist, the remaining $2n$ relations will also hold true.

3. If H be expressed in terms of t and the x 's only, we shall have

$$\frac{dx_i}{dt} = 0; \quad (i = 1, 2, \dots, 2n)$$

and, by integration,

$$x_i = g_i,$$

in which the g 's are arbitrary constants ; whence

$$\frac{dy_i}{dt} = -\frac{\partial H}{\partial g_i}.$$

4. Similarly, if H be expressed in terms of t and the y 's only, we shall have

$$\frac{dy_i}{dt} = 0; \quad (i = 1, 2, \dots, 2n)$$

and, by integration,

$$y_i = g'_i,$$

in which the g 's are arbitrary constants ; whence

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial g'_i}.$$

5. If H be a function of $t, x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n$ only, we shall have

$$\frac{dx_{n+i}}{dt} = 0, \quad \frac{dy_{n+i}}{dt} = 0; \quad (i = 1, 2, \dots, n)$$

and, by integration,

$$x_{n+i} = h_i, \quad y_{n+i} = h'_i, \quad (i = 1, 2, \dots, n)$$

in which the h 's and h 's are arbitrary constants. In this case S may be stated to be a function of t and the n variables x_i and containing n arbitrary constants h_i . The equations

$$y_i = \frac{\partial S}{\partial x_i}, \quad h'_i = \frac{\partial S}{\partial h_i} \quad (i = 1, 2, \dots, n)$$

may then be defined as the general integrals of the simultaneous differential equations

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial y_i}, \quad \frac{dy_i}{dt} = -\frac{\partial H}{\partial x_i} \quad (i = 1, 2, \dots, n)$$

6. If

$$H = (H - R) + R,$$

in which $H - R$ is a function of x_i 's and y_i 's only, while R is expressed in terms of x_{n+i} 's and y_{n+i} 's only; we shall have

$$\left. \begin{aligned} \frac{\partial H}{\partial x_i} &= \frac{\partial (H - R)}{\partial x_i}, & \frac{\partial H}{\partial y_i} &= \frac{\partial (H - R)}{\partial y_i}, \\ \frac{\partial H}{\partial x_{n+i}} &= \frac{\partial R}{\partial x_{n+i}}, & \frac{\partial H}{\partial y_{n+i}} &= \frac{\partial R}{\partial y_{n+i}}; \end{aligned} \right\} (i = 1, 2, \dots, n)$$

whence, if

$$\frac{dx_i}{dt} = \frac{\partial (H - R)}{\partial y_i}, \quad \frac{dy_i}{dt} = -\frac{\partial (H - R)}{\partial x_i},$$

then

$$\frac{dx_{n+i}}{dt} = \frac{\partial R}{\partial y_{n+i}}, \quad \frac{dy_{n+i}}{dt} = -\frac{\partial R}{\partial x_{n+i}};$$

or, employing the idea of variation of constants,

$$\frac{dh_i}{dt} = \frac{\partial R}{\partial h'_i}, \quad \frac{dh'_i}{dt} = -\frac{\partial R}{\partial h_i}.$$

7. If we choose, the x 's and y 's may be divided into two groups containing $2n$ variables each, i. e. n x 's and n y 's, and by solving equations (b) the members of each group may be expressed as functions of the members of the other group only. Suppose such to be the case, and x_i, y_i to be any two members of one of the groups, and x_j, y_j to be any two members of the

remaining group. Then will

$$\frac{\partial y_i}{\partial x_j} = \frac{\partial y_j}{\partial x_i}, \quad (\text{a}') \quad \frac{\partial x_i}{\partial y_j} = \frac{\partial x_j}{\partial y_i}, \quad (\text{c}')$$

$$\frac{\partial y_i}{\partial y_j} = -\frac{\partial x_j}{\partial x_i} \quad (\text{b}') \quad \frac{\partial x_i}{\partial x_j} = -\frac{\partial y_j}{\partial y_i}. \quad (\text{d}')$$

If i' and j' be particular values of i and j , we have from equations (b), remembering that variables belonging to the same group are independent of one another,

$$\frac{\partial y_{i'}}{\partial x_{j'}} = \frac{\partial}{\partial x_{j'}} \frac{\partial S}{\partial x_{i'}} + \sum_i \left[\frac{\partial}{\partial x_i} \frac{\partial S}{\partial x_{i'}} \cdot \frac{\partial x_i}{\partial x_{j'}} \right], \quad (1')$$

$$\frac{\partial y_i}{\partial x_{i'}} = \frac{\partial}{\partial x_{i'}} \frac{\partial S}{\partial x_i} + \sum_j \left[\frac{\partial}{\partial x_j} \frac{\partial S}{\partial x_i} \cdot \frac{\partial x_j}{\partial x_{i'}} \right] = 0;$$

whence

$$\frac{\partial}{\partial x_i} \frac{\partial S}{\partial x_{i'}} = \frac{\partial}{\partial x_{i'}} \frac{\partial S}{\partial x_i} = - \sum_j \left[\frac{\partial}{\partial x_j} \frac{\partial S}{\partial x_i} \cdot \frac{\partial x_j}{\partial x_{i'}} \right]. \quad (2')$$

Substituting (2') in (1'), we have

$$\frac{\partial y_{i'}}{\partial x_{j'}} = \frac{\partial}{\partial x_{j'}} \frac{\partial S}{\partial x_{i'}} - \sum_i \sum_j \left[\frac{\partial}{\partial x_j} \frac{\partial S}{\partial x_i} \cdot \frac{\partial x_j}{\partial x_{i'}} \cdot \frac{\partial x_i}{\partial x_{j'}} \right].$$

Similarly, interchanging i , i' and j , j' ,

$$\frac{\partial y_{j'}}{\partial x_{i'}} = \frac{\partial}{\partial x_{i'}} \frac{\partial S}{\partial x_{j'}} - \sum_j \sum_i \left[\frac{\partial}{\partial x_i} \frac{\partial S}{\partial x_j} \cdot \frac{\partial x_i}{\partial x_{j'}} \cdot \frac{\partial x_j}{\partial x_{i'}} \right];$$

whence

$$\frac{\partial y_{i'}}{\partial x_{j'}} = \frac{\partial y_{j'}}{\partial x_{i'}}; \quad \text{or, in general,} \quad \frac{\partial y_i}{\partial x_j} = \frac{\partial y_j}{\partial x_i}. \quad (\text{a}'')$$

As may easily be seen,

$$\frac{\partial y_i}{\partial x_i} = \sum_j \left[\frac{\partial y_i}{\partial x_j} \frac{\partial x_j}{\partial x_i} + \frac{\partial y_i}{\partial y_j} \frac{\partial y_j}{\partial x_i} \right] = 0, \quad (3')$$

$$\frac{\partial y_j}{\partial x_j} = \sum_i \left[\frac{\partial y_j}{\partial x_i} \frac{\partial x_i}{\partial x_j} + \frac{\partial y_j}{\partial y_i} \frac{\partial y_i}{\partial x_j} \right] = 0, \quad (4')$$

$$\frac{\partial x_i}{\partial y_i} = \sum_j \left[\frac{\partial x_i}{\partial x_j} \frac{\partial x_j}{\partial y_i} + \frac{\partial x_i}{\partial y_j} \frac{\partial y_j}{\partial y_i} \right] = 0, \quad (5')$$

$$\frac{\partial x_j}{\partial y_j} = \sum_i \left[\frac{\partial x_j}{\partial x_i} \frac{\partial x_i}{\partial y_j} + \frac{\partial x_j}{\partial y_i} \frac{\partial y_i}{\partial y_j} \right] = 0. \quad (6')$$

By reason of (a'), (3') and (4') may be written

$$\Sigma_j \frac{\partial y_i}{\partial x_j} \left[\frac{\partial x_j}{\partial x_i} + \frac{\partial y_i}{\partial y_j} \right] = 0 ,$$

$$\Sigma_i \frac{\partial y_i}{\partial x_j} \left[\frac{\partial x_i}{\partial x_j} + \frac{\partial y_j}{\partial y_i} \right] = 0 ;$$

whence (b') and (d') are derived directly by indeterminate coefficients. Also, by reason of (b') and (d'), (5') and (6') may be written

$$\Sigma_j \frac{\partial x_i}{\partial x_j} \left[\frac{\partial x_j}{\partial y_i} - \frac{\partial x_i}{\partial y_j} \right] = 0 ,$$

$$\Sigma_i \frac{\partial x_j}{\partial x_i} \left[\frac{\partial x_j}{\partial y_i} - \frac{\partial x_i}{\partial y_j} \right] = 0 ;$$

from either of which (c') follows immediately by indeterminate coefficients.

UNIVERSITY OF VIRGINIA, *January 24*, 1893.